

## RELATIONSHIP OF PHYSIOLOGICAL VARIABLES TO PLAYING ABILITY OF NATIONAL LEVEL WOMEN HOCKEY PLAYERS

### **Dr. Brij Kishore Prasad**

Associate Prof., Head, Department of Physical Education, Institute of Professional Studies, Gwalior (M.P.).

### **Mr. Sani Kumar Verma**

Asst. Prof., Department of Physical Education, Institute of Professional Studies, Gwalior (M.P.).

### **INTRODUCTION**

The professionals must aware of the latest and high sophisticated tool that can facilitate the measurement of various physiological and morphological characteristic and for different sports event should be given proper extensive training by over prolong period. Physiological variables may be defined as those variables which are directly linked with various physiological systems such as heart rate blood pressure, vital capacity, fat percentage, respiratory rate and hemoglobin. Women's hockey develop separately form men's hockey. Women do not seem to have played hockey widely before the modern era, women hockey was first played at British university and school. The first club, Moberly ladies union club was founded in 1887. Hockey is a national game of India and is very popular in the country, but very less scientific work as been done in the field especially physiological profiles of women hockey players. Research work is very important for advancement of game through which we can educate the coaches, physical education teachers and hockey players regarding the role played by physiological variables in achieving high performance efficiency. Hence the scholar has undertaken the present study.

### **METHODOLOGY**

To conduct the study, Participant (N=25) women hockey players from Madhya Pradesh State Women Hockey Academy, Gwalior were selected as subjects. The entire subjects were residing in the hostel of L.N.I.P.E. Gwalior. The age of subject ranged from 14 to 19 years. On the basis of literature available pertaining to physiological variables of hockey players, opinion of coaches, teachers as well as personal experience of scholar, the following physiological variables were selected for the purpose of the present study. Vital capacity, Breath holding capacity, Peak flow rate, VO<sub>2</sub> max, Resting heart rate and Percentage of fat. Breath holding capacity was measured in seconds by manual breath holding capacity, Vo<sub>2</sub> max for aerobic capacity, was measured in (mls/kg /min) by cooper 12 minute run and walk, Vital capacity (lung volume) was measured in liters by dry spirometer and

Peak flow rate was measured in liter/minute by peak flow meter, Resting heart rate was measured in beats/minute as well as Skinfold thickness measured by applying skin fold caliper on Biceps, triceps, subscapulries and suprailliac sites of the body and recorded nearest to one tenth of a millimeter. The percentage of fat was calculated by referring to assessment of fat percentage table suggested by Durnin and Rahman.

## RESULTS


To examine the relationship of physiological variables with performance of hockey playing ability, Pearson's Product Correlation Moment was used. The level of significance was set at .05, which was considered as adequate for the purpose of the study.

**TABLE 1**

**Relationship of Physiological Variables to Playing Ability in Hockey**

S.NO	Physiological Variables	Correlation Coefficient 'r'
1.	Vital Capacity and Playing ability	.728*
2.	Breath holding capacity and Playing ability	.605*
3.	Peak flow rate and Playing ability	.705*
4.	Vo2 max and Playing ability	.450*
5.	Resting heart rate and Playing ability	-.61*
6.	Percentage of fat and Playing ability	0.04

\* Significant at 0.05 level of significance  $r_{(0.05)(18)} = .444$



## DISCUSSION OF FINDING

Physiological variables of female hockey players of M.P. State Women Hockey Academy and hockey playing ability was found by administering appropriate tests as well as utilization appropriate statistics. It has been found that there is a positive relationship between vital capacity and hockey playing ability. Vital capacity is important for hockey players because the greater vital capacity means greater capacity of individual to inhale air into lungs as well as efficiency of exhale. Oxygen is directly proportional to the amount of air inhaled. The greater amount of oxygen is helpful to a player as adequate oxygen is supplied to the working muscle. Excess of oxygen in the blood delays the onset of lactic acid which may further help players to exhibit performance.

A Significant relationship between breath holding capacity and hockey playing ability may be because it is usually seen that player hold their breath while executing any explosive action or stroke. The holding of breath may assist individual to minimize unwanted movement of the body which may lead to the improvement of the coordination, thereby executing the stroke with greater force as well as accuracy. Further, holding of breath prior execution of strokes may also help an individual to concentrate better.

Peak flow rate denotes the highest rate of expiratory output per minute during highest respiratory function. As and when individual play hockey she is to go for hyper ventilation in order to cope up with highest oxygen demand at that movement. Therefore higher peak flow rate will result in higher ventilation efficiency which will lead to conversation of more oxygen and ultimately

producing more energy. Therefore, probably positive relationship of peak flow rate and playing ability was obtained.

The Significant relationship between VO<sub>2</sub> max and playing ability may be due to the reason that the game of hockey is mainly endurance dominating game and ability of individual to utilize greater oxygen during work. It may significantly help the player to play the game with ease as sufficient oxygen is being utilized by her during contraction of muscles. The finding of the study is supported by M. J. Cosgrove and F.M. Impellizzeri.

A Significant negative relationship between resting heart rate and playing ability may be because resting heart rate and efficiency of heart rate are very closely related. The lower the resting heart rate, the better the efficiency of the heart and this may help in individual to play the game better as hockey is the game where endurance and speed endurance player a very important role. Further, a player may be able to adjust/adopt amicability to the demand of the game as heart rate may not increase to that level where individual may feel uncomfortable.

The Insignificant relationship between percentage of fat and hockey playing ability may be because of the sample selected. As all girls were fairly fit and had been undergoing training last two year. The percentage of fat was optimum in their body. Further keeping the nature in the mind where player play in different position and demand of different position are different. Usually it seen that goalkeeper and fullbacks are taller and well built is in comparison to other players but such body structure shoot position of the player hence insignificant relationship obtained.

## REFERENCE

- N. Apostolidis, "Physiological and Technical Characteristics of Elite Young Basketball Players", *Journal of Sports Medicine Physical Fitness*, Vol. 4 (1), 2004.
- V. Bunc, "Physiological Profile of Very Young Soccer Players", *Journal of Sports Medicine Physical Fitness*, Vol. 3 (2), 2001.
- M. K. Chin, *Physiological Profiles and Sport Specific Fitness of Asian Elite Squash Players*, *British Journal of Sports Medicine* Vol. 12 (2), 1995.
- M. J. Cosgrove, "The Relationship Between Selected Physiological Variables Of Rowers and Rowing Performance as Determined by A 2000 M Ergo-Meter Test", *Journal of Sports Science*. Vol. 3 (1), 1999.
- S. J. Fleck, *Physical and Physiological Characteristics of Elite Women Volleyball Players*", *Canadian Journal Applied Sport Science*., Vol 9 (2), 1985.
- M. R. Green, "Relationship Between Physiological Profiles and On-Ice Performance of a National Collegiate Athletic Association Division I Hockey Team", *Journal of Strength Conditioning and Research*. Vol. 10 (2), 2006.

J. Heller , Physiological Profiles of Male And Female Taekwon-Do (ITF) Black Belts, Journal of Sports Science. Vol. 6 (1), 1998.

F. M. Impellizzeri, “Correlations Between Physiological Variables And Performance In High Level Cross Country Off Road Cyclists”, British Journal of Sports Medicine. Vol. 6 (1), 2005.

L. Leger, “Comparisons Among VO<sub>2</sub> Max Values for Hockey Players and Runners”, Canadian Journal of Applied Sport Science. Vol. 9 (2), 1979.

IJPEHSS